

# Toxicological Effects of Chlorine Dioxide, Chlorite and Chlorate

by Daniel Couri,\* Mohamed S. Abdel-Rahman† and Richard J. Bull‡

Review of the available literature obtained from both acute and chronic experiments utilizing rats, mice and chickens treated with  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  and  $\text{ClO}_3^-$  in drinking water has demonstrated alterations in hematologic parameters in all species tested. The effects were usually dose related and marked changes occurred only at the higher dosages (up to 1000 mg/l.). In chronic studies, rats have been given  $\text{ClO}_2$  at doses of up to 1000 mg/l., and  $\text{NaClO}_2$  or  $\text{NaClO}_3$  at up to 100 mg/l., in their drinking water for one year. Treatment groups receiving  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  or  $\text{ClO}_3^-$  showed alterations in erythrocyte morphology and osmotic fragility; at higher dosages mild hemolytic anemia occurred. An examination of blood glutathione content and RBC enzymes involving glutathione formation showed a dose-related diminution of glutathione in chlorine compound treated groups. The higher oxidative capacity of the chlorine compounds resulting in the decreased erythrocytic glutathione might well be the principal biochemical event leading to the other hematological alterations.

More recent data show that  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  and  $\text{ClO}_3^-$  alter the incorporation of  $^3\text{H}$ -thymidine into the nuclei of various organs of the rat. These data suggest the possibility of increased turnover cells of the gastrointestinal mucosa and inhibited DNA synthesis in several organs. In the latter category, most concern revolves around whether or not the apparent depression of DNA synthesis in the testes is associated with depressed spermatogenesis and reproductive toxicity in the male rat.

## Introduction

Institution of disinfection to drinking water treatment has been one of the key successes of public health policy. In the United States almost complete dependence has been placed on chlorine for primary disinfection. In recent years, however, it has become clear that treatment of drinking water with chlorine results in the formation of trihalomethanes (1, 2). One of the trihalomethanes, chloroform, has been shown to be carcinogenic in mice and rats (3).

These observations stimulated a search for a means to minimize the formation of these products. One of the most attractive alternatives is to substitute disinfection methods that do not promote the formation of chlorinated by-products.

Chlorine dioxide is a very attractive alternative to chlorine as a disinfectant. For all intents and purposes its disinfectant properties are equivalent to or perhaps exceed those of chlorine (4). Chlorine dioxide does not react with phenol to produce the same taste and odor problems that result from chlorine treatment (5). It has the added advantage that formation of trihalomethanes does not occur (6, 7).

Prior to the recent consideration of chlorine dioxide as a primary disinfectant for potable water, very little information existed concerning its toxic-

\*Department of Pharmacology, College of Medicine, Ohio State University, Columbus, Ohio 43210.

†Department of Pharmacology, CMDNJ-New Jersey Medical School, Newark, New Jersey 07103.

‡Health Effects Research Laboratory, U.S. Environmental Protection Agency, Cincinnati, Ohio 45268.

cology. The use of chlorine dioxide for this purpose also requires consideration of by-products of its use that will occur in the consumed product, chlorite ( $\text{ClO}_2^-$ ) and chlorate ( $\text{ClO}_3^-$ ). Chlorite occurs both as a result of being the first reduction product in oxidative reactions in which  $\text{ClO}_2$  participates and because  $\text{ClO}_2$  is often generated by acidifying a sodium chlorite feed solution. Chlorate on the other hand arises primarily by a disproportionation reaction of  $\text{ClO}_2$  that is catalyzed by ultraviolet light (8), giving rise to one molecule of  $\text{ClO}_2^-$  and one molecule of  $\text{ClO}_3^-$  per two molecules of  $\text{ClO}_2$ .

As a result of the interest in  $\text{ClO}_2$  as a primary disinfectant, a number of studies have been initiated to investigate the toxicity associated with  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  and  $\text{ClO}_3^-$ . This paper reviews the literature which has resulted from these investigations and attempts to deal with the mechanisms involved with the cellular damage induced by these agents.

## Hematologic Effects

Rats, mice and chickens have been treated with  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  and/or  $\text{ClO}_3^-$  in drinking water at concentrations ranging from 1 to as high as 1000 mg/l. A relatively consistent picture has been observed across species. In the rat, Heffernan et al. (9) demonstrated that chlorite concentrations of 100 mg/l. and above resulted in decreased red blood cell counts, hemoglobin concentrations and packed cell volume at 30 and 60 days of exposure. These effects tended to be reversed by 90 days of exposure despite continued treatment with sodium chlorite. These data were extended to the mouse by Moore and Calabrese (10) with very similar results. Abdel-Rahman et al. (11) studied the effects of  $\text{ClO}_2$  and  $\text{ClO}_3^-$  as well as  $\text{ClO}_2^-$  in both the rat and chicken for periods of up to 9 months. Signs of anemia observed in shorter term exposures of 30 to 60 days (12) had substantially disappeared in this interval with  $\text{ClO}_2$  and  $\text{ClO}_2^-$ . However, the results obtained with  $\text{ClO}_3^-$  at 9 months indicated 13 and 28% decreases in RBC at 10 and 100 mg/l. This represented a considerably greater effect than had been observed at shorter time intervals.

Accompanying the relatively small decreases in RBC count is subclinical evidence of increased red cell destruction. This was first reported by Heffernan et al. (9) in the cat, where significant decreases in the half-life of erythrocytes were observed at concentrations of  $\text{ClO}_2^-$  of 100 mg/l. of drinking water and above. No effect was observed at 10 mg  $\text{ClO}_2^-$ /l. Other investigators have provided evidence of distortions in erythrocyte morphology in mice (10) as well as the rat and chicken (12). These

changes were quite marked at concentrations of 100 mg  $\text{ClO}_2^-$ /l. ranging from mildly crenated erythrocytes which maintained their discoid shape to crenated spheres (echinocytes). With  $\text{ClO}_3^-$  treatment, codocytes (Mexican hat-shaped) were also observed in the rat. In the chicken, the characteristic abnormality was the dacrocyte, which was evident primarily with  $\text{ClO}_2$  treatments of 100 and 1000 mg/l. Early studies had somewhat paradoxically indicated that  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  and  $\text{ClO}_3^-$  decreased osmotic fragility of erythrocytes isolated from treated animals (12). Subsequent experiments have shown this effect is most likely due to the oxidation of thiol groups resulting in the precipitation of hemoglobin to give falsely low indications of hemolysis (13). This is also consistent with the *in vitro* observations of Heffernan et al. (14) that a complete loss of measurable pigment (both hemoglobin and methemoglobin) occurred at high concentrations of  $\text{ClO}_2^-$ .

## Biochemical Effects

Early reports had documented the ability of  $\text{ClO}_2^-$  to oxidize hemoglobin to methemoglobin by high intraperitoneal doses (320 mg/kg) (15).

Heffernan et al. (9) confirmed these observations using intraperitoneal doses in the rat and bolus oral doses of  $\text{ClO}_2^-$  in the cat. However, in studies where  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  or  $\text{ClO}_3^-$  was included in the drinking water for up to several months, no significant increases in methemoglobin concentrations resulted with doses as high as 1000 mg/l. in the rat, mouse or chicken (9, 10, 12).

Therefore, although a real hazard with large bolus doses, methemoglobinemia appears to be an unlikely hazard at concentrations of  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  and  $\text{ClO}_3^-$  which might be anticipated to be associated with drinking water treatment.

Heffernan et al. (9) reported that  $\text{ClO}_2^-$  incubated with RBC *in vitro* depleted these cells of glutathione and increased the level of hydrogen peroxide produced as measured by catalase complex 1 formation. These *in vitro* studies clearly demonstrated that virtually complete loss of glutathione occurred before there was significant accumulation of methemoglobin. Additionally, these studies demonstrated a biphasic consumption of added  $\text{ClO}_2^-$ , one phase of which was associated with hydrogen peroxide generation and glutathione depletion and the other with hemoglobin oxidation. The changes in erythrocyte morphology by scanning electron microscopy *in vitro* paralleled exactly the progression observed with  $\text{ClO}_2$  treatment in subsequent *in vivo* experiments (10, 12).

From these data it was clear that  $\text{ClO}_2^-$  possesses

the types of activity associated with chemicals that produce oxidative hemolysis *in vivo* (16–18).

Red blood cell glutathione concentrations have proven to be the most sensitive measure of  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  and  $\text{ClO}_3^-$  induced hematological effects *in vivo* as well. Heffernan et al. (9) observed significant depletions of glutathione and increases of 2,3-diphosphoglycerate in the red cells of rats exposed to as little as 50 mg of  $\text{ClO}_2^-$ /l. of drinking water. Unlike the effects on RBC counts and hemoglobin concentrations, tolerance does not develop to  $\text{ClO}_2^-$ -induced depletion of glutathione. Abdel-Rahman et al. (11, 12), confirmed this result with  $\text{ClO}_2^-$  and extended the observations to  $\text{ClO}_2$  and  $\text{ClO}_3^-$ . Like  $\text{ClO}_2^-$ ,  $\text{ClO}_3^-$  produced stable depression of red cell glutathione concentrations with continued treatment at 10 and 100 mg/l. On the other hand, depression of red cell glutathione levels by  $\text{ClO}_2$  tends to disappear with continued exposure (11, 13).

The recovery or nonrecovery of glutathione levels with continued treatment *in vivo* appeared at least partially dependent upon a complex set of interactions with redox systems operating in the rat erythrocyte (19). The reversibility of the  $\text{ClO}_2$  effect appears to result from increased levels of glutathione reductase activity in the red cell following 6 months of treatment. In the case of  $\text{ClO}_3^-$  there appears to be no such adaptive increase in this enzyme level. With  $\text{ClO}_2^-$  the glutathione reductase levels do increase to levels comparable to those observed with  $\text{ClO}_2$ . However, this adaptive change appears to be offset by significantly depressed levels of catalase in the red cell with  $\text{ClO}_2^-$  treatment.

The importance of the changes in enzyme levels in the adaptation to chronic  $\text{ClO}_2$  exposure has been indirectly supported by the data of Heffernan et al. (9, 14). First, the addition of glucose to red cells treated with  $\text{ClO}_2^-$  *in vitro* partially protects against depletion of glutathione. Glutathione peroxidase is the principal means of disposing of hydrogen peroxide in the red cell (17). Glucose serves as the carbon source which donates the reducing equivalents necessary for the reduction of oxidized glutathione (17, 20).

Secondly, *in vivo* treatment with  $\text{ClO}_2^-$  greatly enhances the formation of  $\text{ClO}_2^-$ -induced formation of hydrogen peroxide *in vitro*. Clearly *in vivo* treatment with  $\text{ClO}_2^-$  reduces the ability of the red cell to dispose of hydrogen peroxide. Although the animal is not seriously ill at levels of  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  or  $\text{ClO}_3^-$  considerably exceeding those contemplated in drinking water treatment, the body's ability to adapt to oxidant stress is compromised at relatively low concentrations of these chemicals.

These considerations focus attention on populations known to be sensitive to oxidants, for example, those deficient in glucose-6-phosphate dehydrogenase. Such individuals are known to be considerably more sensitive to oxidant stress than normal humans. As will be discussed in other papers in this symposium, erythrocytes isolated from such individuals are three to four times as susceptible to  $\text{ClO}_2^-$ -induced damage than normal human red cells (21). In this regard, Michael et al. (22) conducted a human study in a small town which utilized  $\text{ClO}_2$  in combination with chlorine as a primary disinfectant during the summer months. Concentrations of  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  and  $\text{ClO}_3^-$  were not closely controlled but the total of their concentrations averaged 5–7 mg/l. In this situation normal humans were not apparently affected. However, the only individual in the community deficient in glucose-6-phosphate dehydrogenase experienced a substantial decrease in hemoglobin levels and hematocrit. With only one case it is difficult to conclusively ascribe this change to  $\text{ClO}_2$  or its by-products, but it does suggest reason for concern about sensitive populations.

## Evidence of Damage to Other Tissues

Research examining nonhematologic effects of  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  and  $\text{ClO}_3^-$  is sparse. Distribution and clearance of  $^{36}\text{Cl}$  derived from  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  or  $\text{ClO}_3^-$  is widespread throughout all organs including bone marrow (12, 23, 24).

There is some evidence of macromolecular binding in the liver. These findings raised the question of whether or not cellular types in addition to the red cell are similarly targeted by these chemicals.

The gastrointestinal mucosa is the obvious first potential target of residual disinfectant species. Abdel-Rahman et al. (11) examined the incorporation of  $^3\text{H}$ -thymidine into nuclei of cells of the intestinal mucosa of the rat as a measure of the turnover of these cells. At levels of 10 mg/l. of  $\text{ClO}_2$  and  $\text{ClO}_2^-$  in drinking water for 3 months, there were significant increases in  $^3\text{H}$ -thymidine incorporation, implying an increased turnover of intestinal epithelium. No effect was observed with equivalent concentrations of  $\text{ClO}_3^-$ . Unfortunately, lower levels of exposure to  $\text{ClO}_2$  and  $\text{ClO}_2^-$  were not examined.

Examination of other organs revealed that  $\text{ClO}_2$  treatment inhibited  $^3\text{H}$ -thymidine incorporation into nuclei of cells in the kidney and testes of the rat.  $\text{ClO}_2^-$  produced similar effects in the testes and markedly inhibited  $^3\text{H}$ -thymidine incorporation in the liver at 10 and 100 mg/l.  $\text{ClO}_3^-$  produced significant effects only in the testes. At 100 mg/l,  $\text{ClO}_2^-$  also

inhibited the incorporation of  $^3\text{H}$ -thymidine into the gastrointestinal mucosa and  $\text{ClO}_2$  produced a smaller increase than had been observed with 10 mg/l. These data suggest that at high doses in the intestinal mucosa and at lower doses in testes and liver,  $\text{ClO}_2$  and/or its by-products is interfering with DNA synthesis. The significance of these results is presently difficult to determine. But since a large portion of the DNA synthesis ongoing in testes is associated with spermatogenesis, these results cannot be lightly dismissed. Consequently, it is essential that this work be followed up with studies of the reproductive toxicity of these chemicals and be extended to lower doses.

## Conclusions

It is clear from this brief review that substantial questions remain to be answered concerning the safety of  $\text{ClO}_2$  as a primary disinfectant of drinking water. Although  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  and  $\text{ClO}_3^-$  have been used extensively in drinking water treatment, actual human exposure has been limited in the past because the concentrations proposed for taste and odor control are much lower than required for primary disinfection. Europeans have used  $\text{ClO}_2$  as a primary disinfectant, but again levels which have been used are quite low because disinfectants are employed after much of the organic material has been removed from the water by granular activated carbon, a practice not common in the U.S. Consequently, only in certain very limited circumstances have epidemiological evaluations been possible and appropriate to the use of  $\text{ClO}_2$  as a primary disinfectant. Consequently, it appears that further experimental work will be necessary to more clearly identify and quantify the hazards involved with its use.

Three major areas of concern over the use of  $\text{ClO}_2$  as a disinfectant have been identified by research performed to date. First, there is a need to study human populations which display higher levels of sensitivity to oxidant chemicals. Second, efforts must be made to determine the significance of increased turnover of the epithelium of the gastrointestinal tracts at low doses of  $\text{ClO}_2$  and  $\text{ClO}_2^-$ . This tissue normally turns over quite rapidly. On the other hand, cellular damage followed by regeneration has been associated with increased susceptibility to chemical carcinogens in other organs such as the liver (25). Third, and perhaps most important, is the question of potential reproductive effects of  $\text{ClO}_2$  and its by-products implied, but not established, by the reduced incorporation of  $^3\text{H}$ -thymidine into testicular DNA. The fact that this effect occurred at quite low doses of  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  and  $\text{ClO}_3^-$

raises the level of concern considerably. It should be noted that similar effects were not observed with  $\text{HOCl}$  (11).

In summary, there is little conclusive evidence to prevent the use of  $\text{ClO}_2$  as a disinfectant at present. On the other hand, there are scattered pieces of indirect evidence that would advise caution about instituting the use of  $\text{ClO}_2$  as a primary disinfectant in place of chlorine. This is complicated by the fact that the total concentration of  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  and  $\text{ClO}_3^-$  might reach several parts per million in the absence of removing background organic material to be an effective disinfectant. It is obvious that this question can only be finally resolved through further research.

## REFERENCES

1. Rook, J. J. Formation of haloforms during chlorination of natural waters. *J. Water Treat. Exam.* 23: 234-236 (1974).
2. Bellar, T. A., Lichtenberg, J. J., and Kroner, R. D. The occurrence of organohalides in chlorinated drinking water. *J. Am. Water Works Assoc.* 66: 703-706 (1974).
3. NCI. Report on the Carcinogenesis Bioassay of Chloroform, National Cancer Institute, 1976, National Technical Information Service, Springfield, Va., PB-264018.
4. Akin, E. W., Hoff, J. C., and Lippy, E. C. Waterborne outbreak control: which disinfectant? *Environ. Health Perspect.* 46: 7-12 (1982).
5. Enger, M. Treatment of water with chlorine dioxide to improve the taste. *Gass Wasser Fach* 14: 330-336 (1960).
6. Stevens, A., Seeger, D., and Slocum, C. J. Products of chlorine dioxide treatment of organic material in water. Paper presented at Workshop on Ozone and Chlorine Dioxide Oxidation Products of Organic Materials, Cincinnati, Ohio, November 17-19, 1976.
7. Miltner, R. The effect of chlorine dioxide on trihalomethanes in drinking water. Master Thesis, University of Cincinnati, 1976, pp. 20-50.
8. Stevens, A. A. Reaction products of chlorine dioxide. *Environ. Health Perspect.* 46: 101-110 (1982).
9. Heffernan, W. P., Guion, C., and Bull, R. J. Oxidative damage to the erythrocyte induced by sodium chlorite, *in vivo*. *J. Environ. Pathol. Toxicol.* 2: 1487-1499 (1979).
10. Moore, G. S. and Calabrese, E. J. The effects of chlorine dioxide and sodium chlorite on erythrocytes of A/J and C57L/J mice. *J. Environ. Pathol. Toxicol.* 4(2-3): 513-524 (1980).
11. Abdel-Rahman, M. S., Couri, D., and Bull, R. J. Toxicity of chlorine dioxide in drinking water. *J. Environ. Pathol. Toxicol.*, in press.
12. Abdel-Rahman, M. S., Couri, D., and Bull, R. J. Kinetics of  $\text{ClO}_2$  and effect of  $\text{ClO}_2$ ,  $\text{ClO}_2^-$  and  $\text{ClO}_3^-$  in drinking water on blood glutathione and hemolysis in rat and chicken. *J. Environ. Pathol. Toxicol.* 3: 431-449 (1980).
13. Abdel-Rahman, M. S., Couri, D., and Bull, R. J. Effect of exogenous glutathione, glutathione reductase, chlorine dioxide and chlorite on osmotic fragility of rat blood *in vitro*. *J. Environ. Pathol. Toxicol.*, in press.
14. Heffernan, W. P., Guion, C., and Bull, R. J. Oxidative damage to the erythrocyte induced by sodium chlorite, *in vitro*. *J. Environ. Pathol. Toxicol.* 2: 1501-1510 (1979).
15. Musil, J., Kontek, Z., Chalupa, J., and Schmidt, P. Toxicological aspects of chlorine dioxide application for the treat-

- ment of water containing phenol. *Chem.-Technol. Praha* 8: 327-345 (1964).
16. Allen, D., and Jandle, J. Oxidative hemolysis and precipitation of hemoglobin. *J. Clin. Invest.* 39: 1818-1836 (1960).
  17. Cohen, G., and Hochstein, P. Generation of hydrogen peroxide in erythrocytes by hemolytic agents. *Biochemistry* 3: 895-900 (1964).
  18. Kiese, M. *Methemoglobinemia: A Comprehensive Treatise*. CRC Press, Cleveland, 1974.
  19. Couri, D., and Abdel-Rahman, M. S. Effect of chlorine dioxide and metabolites on glutathione dependent system in rat, mouse and chicken blood. *J. Environ. Pathol. Toxicol.* 3: 451-460 (1979).
  20. Jacobs, H. S. and Jandl, J. H. Effects of sulfhydryl inhibition on red blood cells. III. Glutathione in the regulation of the hexose monophosphate pathway. *J. Biol. Chem.* 241: 4243-4250 (1966).
  21. Moore, G. S., Calabrese, E. J. and Ho, S. C. Groups at potentially high risk from chlorine dioxide treated water. *J. Environ. Pathol. Toxicol.* 4 (2,3): 465-470 (1980).
  22. Michael, G. E., Miday, R. K., Berez, J. P., Miller, R. G., Greathouse, D. G., Kraemer, D. F. and Lucas, J. B. Chlorine dioxide water disinfection: a prospective epidemiology study. *Arch. Environ. Health* 36: 20-27 (1981).
  23. Abdel-Rahman, M. S., Couri, D., and Jones, J. D. Chlorine dioxide metabolism in rat. *J. Environ. Pathol. Toxicol.* 3: 421-430 (1980).
  24. Abdel-Rahman, M. S., Couri, D., and Bull, R. J. The kinetics of chlorite and chlorate in rat. *J. Environ. Pathol. Toxicol.*, in press.
  25. Pound, A. W., and McGuire, L. J. Repeated partial hepatectomy as a promoting stimulus for carcinogenic response of liver to nitrosamines in rats. *Brit. J. Cancer* 37: 585-594 (1978).